

ABOUT BECKMAN

BIOLOGICAL INTELLIGENCE

HUMAN-COMPUTER
INTELLIGENT INTERACTION

INTEGRATIVE IMAGING

MOLECULAR AND ELECTRONIC
NANOSTRUCTURES

STRATEGIC INITIATIVES

DEVELOPING INITIATIVE

FACULTY

FELLOWS

FACILITIES

NEWS AND EVENTS

News Archive

Synergy

News by Topic

Event Calendar

Director's Seminar

Thursdays at 12:20

Space Reservations

Beckman at 20

SERVICES

GIVING


DIRECTORY

Search

RSS: News | Events | Café Menu

FOLLOW US ON JOIN OUR GROUP 

New Dipstick Method for Detecting Lead in Paint Reported



Published February 17, 2010

By Steve McGaughey, Beckman Institute Writer



Yi Lu

Beckman Institute researcher **Yi Lu** took another researcher's discovery about a new way to achieve catalytic reactions and turned it on its head – leading to a completely new research line for him and a simple, low-cost lead paint detection method for home and official use.

In 1994 Lu read in *Chemistry and Biology* magazine a seminal work by Ronald Breaker and Gerald Joyce about a new method of using metal ions to turn DNA into an enzyme to catalyze chemical reactions – and wondered what would happen if the process were reversed.

"We asked 'can we actually turn DNAzymes that bind to a metal ion like lead or mercury into "reporters" for the presence of metal ions?" said Lu, a researcher at Illinois's Beckman Institute.

So Lu leveraged his laboratory's previous work on how different metal ions can help DNA function and flipped the direction to see if the DNA enzymes, or DNAzymes (deoxyribozymes) as they are called, can help detect metal ions. The research line proved so successful that it led to the creation of dipstick test prototypes for detecting toxic metals.

In a January paper in *Chemical Communications* titled *Easy-to-use dipstick tests for detection of lead in paints using non-cross-linked gold nanoparticle-DNAzyme conjugates*, Lu and his collaborators report that they have fashioned a viable, easy-to-use dipstick test that, unlike other methods, requires no instrumentation or laboratory work and that meets EPA guidelines for lead detection.

The Lu group had previously reported using cross-linked gold nanoparticles and aptamers, DNAs that bind many targets, in a lateral flow device – also known as a dipstick test and similar to commonly used home pregnancy tests – to create colorimetric biosensors for detecting molecules such as cocaine (see *Angew. Chem., Int. Ed.*, 2006, 45, 7955-7959). The biosensors were highly sensitive and accurate, but it was difficult to apply the method to DNAzyme-based sensing of metal ions due to the difficulty in controlling DNAzyme catalytic activity and product release.

This recent paper reported on an advancement in which the researchers used non-cross-link DNAzyme-gold nanoparticle conjugates to create a hybridization method for detecting lead that is stable at ambient conditions for long periods of time. The sensor is able to detect trace levels of lead that more than meets EPA requirements.

The dipstick test shows a color change to red when lead is detected at one milligram per square centimeter (the level paint is deemed to be lead-based), and has been shown to be effective even if the lead-based paint has been repeatedly painted over.

The researchers write that "Designing easy-to-use biosensors for trace metal ions in the environment is of considerable importance as these metal ions are large in number, small in quantity and high in toxicity."

They add that their DNAzyme-gold nanoparticle non-cross-link method, as applied in a dipstick test, "will find wide use in household and other environmental applications."

To read the paper, click [here](#).

 SHARE

 PRINT