


ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

Vibrational Dynamics of Oxygenated Heme Proteins[†]

View Online

Wei-qiao Zeng,^a Alexander Barabanschikov,^a Ningyan Wang,^b Yi Lu,^b Jiyong Zhao,^c Wolfgang Sturhahn,^c E. Ercan Alp^c, and J. Timothy Sage^{*a}

Received Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

First published on the web Xth XXXXXXXXXX 200X

DOI: 10.1039/b000000x

Advanced spectroscopic techniques coupled with DFT calculations reveal the vibrational dynamics of the iron in stable dioxygen complexes with myoglobin and with a mutant engineered to model the catalytic site of heme-copper oxygenases. The unprecedented level of detail will constrain computational modelling of reactions with oxygen.

Heme proteins dominate biological O₂ consumption on Earth. The reduction of molecular oxygen to water by heme-copper oxidases (HCOs) powers the metabolism of aerobic organisms,¹ and numerous heme proteins function as O₂ sensors and transporters.² Globins are heme proteins employed throughout the tree of life³ as O₂ sensors and enzymes, but are most familiar for their roles in O₂ transport and storage in vertebrate metabolism.⁴ Heme-O₂ complexes (Fig. 1) hold great interest as intermediates in diverse enzymatic processes, but detection and interpretation of their vibrational properties remain challenging.

Nuclear resonance vibrational spectroscopy (NRVS; also known as nuclear inelastic scattering) provides a comprehensive and quantitative picture of the vibrational dynamics of ⁵⁷Fe in isotopically enriched proteins.⁶ Fig. 2a and b compares vibrational densities of states (VDOS) for the heme iron atom derived from NRVS measurements on frozen solutions of O₂ complexes with the muscle pigment myoglobin (Mb) and with a L29H/F43H mutant (Cu_BMb), engineered to coordinate a second metal ion via histidines at positions 29, 43, and 64. With Cu(I) occupying the engineered site, Cu_BMb mimics the structure (Fig. S1[†]) of HCOs, and achieves partial reduction of O₂ to produce peroxide.⁷ On the other hand, substitution of the redox inert Ag(I) ion for Cu in air yields a stable Ag(I)–Cu_BMbO₂ complex,⁸ which models structural




Fig. 1 Crystallographic model of the heme site of oxygenated myoglobin refined to 1.25 Å (PDB code 2Z6S),⁵ with spheres highlighting the heme iron and O₂ ligand.

and electrostatic features of the first reaction intermediate.

The Fe dynamics of the two complexes (Fig. 2) exhibit both qualitative similarities and quantitative differences. Features are clearly resolved from the Ag(I)–Cu_BMbO₂ VDOS at approximately 339, 370, 428 and 588 cm⁻¹. The latter two appear at higher frequencies than corresponding features in the MbO₂ VDOS, reflecting the influence of the second metal ion on the oxygenated heme. Although we are not aware of previous applications of NRVS to oxygenated heme proteins, recent measurements on oxygenated “picket fence” porphyrins⁹ identified vibrations of the FeOO fragment within 10 cm⁻¹ of the 423 and 579 cm⁻¹ features observed for MbO₂.

Isotopic labelling also distinguishes Fe vibrations from the dominant signal due to porphyrin vibrations in Raman scattering from frozen solutions of these molecules (Fig. 3). In particular, the 2.3 cm⁻¹ upshift of the 579 cm⁻¹ MbO₂ frequency when ⁵⁴Fe replaces ⁵⁷Fe is consistent with the 2 cm⁻¹

[†] Electronic Supplementary Information (ESI) available: methods, supplementary tables and figures, and animations of predicted 215, 298, 346, 373, 449, and 639 cm⁻¹ vibrations. See DOI: 10.1039/b000000x/

^a Department of Physics and Center for Interdisciplinary Research on Complex Systems, Northeastern University, Boston, MA 02115. E-mail: jt-sage@neu.edu

^b Department of Chemistry, University of Illinois at Urbana Champaign, Urbana, IL 61801.

^c Advanced Photon Source, Argonne National Laboratory, Argonne, IL 60439.




Fig. 2 The Fe VDOS determined from NRVS measurements on (a) Ag(I)–Cu_BMbO₂ and (b) horse MbO₂ completely describe vibrational motion of Fe in each molecule. Shading indicates areas determined by curve fitting the VDOS, in cyan, and estimated from ⁵⁷Fe/⁵⁴Fe isotope shifts, in magenta. The black trace in panel (c) shows the predicted Fe VDOS, indicating regions where each Fe-ligand bond contributes to the signal. Predicted contributions from motion along *x* (red), *y* (magenta), and *z* (blue) refer to the coordinate system in Fig. 1 where the four pyrrole nitrogens define the *xy* plane and the FeOO group defines the *xz* plane.

shift predicted (by Eq. S10) based on the 0.11 area of the 579 cm⁻¹ feature in the Fe VDOS. This frequency drops to 572 cm⁻¹ at ambient temperature (Fig. S2), where previous Raman measurements using O₂ isotopomers¹⁰ reveal frequency shifts that we use in conjunction with the NRVS results to experimentally quantify the kinetic energy distribution (KED). More than 90% of the mode energy is localized on the FeOO fragment, with the majority (77%) on one of the oxygen atoms. The asymmetric KED is an unambiguous signature for end-on binding and distinct from the more uniform KED predicted for a two-body Fe–O₂ oscillator (Table 1).

In contrast with ambient temperature Raman measurements (Fig. S2), improved spectral resolution at cryogenic temperatures also identifies an Fe-isotope sensitive peak at 367 cm⁻¹ in MbO₂ that approximately corresponds in both frequency and predicted area (using Eq. S10) to the 369 cm⁻¹ NRVS feature, as indicated by the shaded area in Fig. 2b. Subtracting the ⁵⁷Fe and ⁵⁴Fe-enriched MbO₂ Raman spectra (Fig. 3) highlights the shifts of the 579 and 367 cm⁻¹ bands. Ad-




Fig. 3 Cryogenic Raman spectra recorded in resonance with the Soret absorption band include contributions from Fe vibrations, highlighted by the ⁵⁷Fe/⁵⁴Fe difference spectrum, among more strongly enhanced porphyrin vibrations.

ditional differences in the 340–380 cm⁻¹ region and near 424 cm⁻¹ are difficult to quantify because they may not correspond to simple frequency shifts, although their proximity to strong NRVS signals is suggestive.

DFT calculations on a model for MbO₂ (Fig. S3) yield the VDOS for Fe (Fig. 2c), as well as for other atoms (Figs. S4, S5). Significant FeOO motion contributes to a predicted 639 cm⁻¹ mode (Movie S639[†]), which exhibits a KED centered on the O_b atom bound to the heme Fe in agreement with the observed 579 cm⁻¹ mode KED (Table 1, Fig. S4), and an unresolved pair of modes with predicted 449 (Movie S449[†]) and 451 cm⁻¹ frequencies, possessing a total $e_{\text{Fe}}^2 = 0.42$, accounting for much of the 423 cm⁻¹ VDOS area. An additional predicted mode at 298 cm⁻¹ (Movie S298[†]) with 30% localization on FeOO does not correspond to a clearly resolved feature in the Fe VDOS.

The predicted 375 cm⁻¹ vibration (Movie S375[†]) corresponds closely in frequency and Fe amplitude to the band observed near 370 cm⁻¹ in both NRVS and Raman. Vibrational correlation analysis¹² reveals 34% overlap with the γ_7 mode of iron porphyrin, which involves Fe motion perpendicular to the heme plane coupled with concurrent methine displacements, and visual examination also reveals in-plane Fe motion, coupled with rotation of the His64 imidazole ring.

Fe motion parallel to the heme plane contributes throughout the 250–400 cm⁻¹ region, distinctly partitioned either parallel or perpendicular to the FeOO plane, similar in character to

Table 1 Experimental vibrational kinetic energy distributions over the FeOO group

protein	frequency (cm ⁻¹)	e_{Fe}^2	$e_{\text{O}_b}^2$	$e_{\text{O}_b}^2$
Ag(I)-Cu _B MbO ₂	428 ^a	0.52 ^a	–	–
Ag(I)-Cu _B MbO ₂	588 ^a	0.08 ^a	–	–
MbO ₂	423 ^a	0.52 ^a	–	–
MbO ₂	579 ^a	0.11 ^a	0.77 ^b	0.06 ^b
MbO ₂ model ^c	298	0.13	0.04	0.11
MbO ₂ model ^c	449/451	0.42	0.09	0.17
MbO ₂ model ^c	639	0.13	0.63	0.03
Fe-O ₂ oscillator	–	0.36	0.34	0.34

^a Determined from cryogenic NRVS measurements. ^b Determined according to Eq. S10 from isotopic frequency shifts reported at ambient temperature.^{10,11} ^c DFT calculation on optimized model for the MbO₂ heme site.

the vibrational anisotropy observed experimentally for single crystals of Fe nitrosyl porphyrins.¹³ Specifically, comparison with the predicted VDOS implicates Fe motion perpendicular to the FeOO plane as the primary contribution to the sharp maximum in the observed signal at 337 cm⁻¹, while vibrations with significant Fe motion parallel to the FeOO plane are distributed over a wider range of frequencies.

The correspondence between predicted and observed features becomes less apparent at lower frequencies. The active site model is too small to capture global oscillations of the polypeptide that carry the heme group and account for the broad feature peaked near 30 cm⁻¹. It is conceivable that coupling with polypeptide vibrations also accounts for failure to resolve the predicted 215 cm⁻¹ feature (Movie S215[†]), although NRVS measurements on oxygenated picket fence porphyrins reveal a feature attributed to Fe-imidazole stretching near 200 cm⁻¹.⁹

Reproduction of the areas of prominent features above 300 cm⁻¹ suggests that the MbO₂ model calculation reproduces the character of the corresponding vibrations. However, the closed-shell singlet state calculation overestimates FeOO vibrational frequencies (Table 1), mirroring results on picket fence porphyrins, which found that spin-unrestricted DFT calculations predict an open-shell electronic configuration that significantly underestimates these frequencies.⁹ Quantitative vibrational comparisons should inform the ongoing discussion of the electronic ground state of oxygenated hemes.¹⁴

Although 1 atmosphere of CO is not sufficient to saturate the heme site of Cu(I)-Cu_BMb,¹⁵ Ag(I)-Cu_BMb is fully saturated with O₂ when prepared in air.⁷ This suggests that O₂ affinity exceeds CO affinity by at least an order of magnitude, in contrast with native Mb, whose affinity for CO is 25 times larger than for O₂.¹⁶ The enhanced discrimination in favor of the O₂ substrate by Ag(I)-MbO₂ is notable, since His64 ligates Ag(I) and is no longer available to hydrogen bond with

O₂, as in MbO₂ (Fig. 1). Apparently, the Ag(I) ion more than compensates for the loss of the stabilizing hydrogen bond.

The increased FeOO frequencies observed for Ag(I)-Cu_BMbO₂ in comparison with MbO₂ are consistent with a strengthened Fe-O₂ bond. The Fe centered 428 cm⁻¹ FeOO vibration of Ag(I)-Cu_BMbO₂ presumably corresponds to the 435 cm⁻¹ frequency reported in the transiently formed O₂ complex with cytochrome *c* oxidase,¹⁰ and both FeOO frequencies lie at the upper end of the range reported for oxygenated heme proteins (Table S1). Thus, Cu_BMb reproduces vibrational as well as functional HCO properties. Protein engineering yields a stable O₂ complex that mimics the first intermediate in the reaction of fully reduced HCO with O₂, and its vibrational behavior should guide the course of computational investigations.

We acknowledge financial support from the National Science Foundation (CHE-1026369) and the National Institutes of Health (GM-062211). Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Basic Energy Sciences, Office of Science, under Contract No. DEAC02-06CH11357.

References

- 1 P. Brzezinski and R. B. Gennis, *J Bioenerg Biomembr*, 2008, **40**, 521–531.
- 2 K. R. Rodgers, *Curr Opin Chem Biol*, 1999, **3**, 158–167.
- 3 S. N. Vinogradov and L. Moens, *J Biol Chem*, 2008, **283**, 8773–8777.
- 4 J. B. Wittenberg, *Gene*, 2007, **398**, 156–161.
- 5 M. Unno, H. Chen, S. Kusama, S. Shaik and M. Ikeda-Saito, *J Am Chem Soc*, 2007, **129**, 13394–13395.
- 6 W. Zeng, N. J. Silvermail, W. R. Scheidt and J. T. Sage, in *Nuclear Resonance Vibrational Spectroscopy (NRVS)*, John Wiley & Sons, Ltd., Chichester, UK, 2007, pp. 401–421.
- 7 J. A. Sigman, H. K. Kim, X. Zhao, J. R. Carey and Y. Lu, *Proc Natl Acad Sci USA*, 2003, **100**, 3629–3634.
- 8 J. A. Sigman, B. C. Kwok and Y. Lu, *J. Am. Chem. Soc.*, 2000, **122**, 8192–8196.
- 9 J. Li, Q. Peng, A. Barabanshikov, J. W. Pavlik, E. E. Alp, W. Sturhahn, J. Zhao, C. E. Schulz, J. T. Sage and W. R. Scheidt, *Chemistry*, 2011, **17**, 11178–11185.
- 10 S. Hirota, T. Ogura, E. H. Appelman, K. Shinzawah-Itoh, S. Yoshikawa and T. Kitagawa, *J. Am. Chem. Soc.*, 1994, **116**, 10564–10570.
- 11 T. Tomita, S. Hirota, T. Ogura, J. S. Olson and T. Kitagawa, *J. Phys. Chem. B*, 1999, **103**, 7044–7054.
- 12 A. Barabanshikov, A. Demidov, M. Kubo, P. M. Champion, J. T. Sage, J. Zhao, W. Sturhahn and E. E. Alp, *J Chem Phys*, 2011, **135**, 015101.
- 13 J. W. Pavlik, A. Barabanshikov, A. G. Oliver, E. E. Alp, W. Sturhahn, J. Zhao, J. T. Sage and W. R. Scheidt, *Angew. Chem. Int. Ed.*, 2010, **49**, 4440–4404.
- 14 H. Chen, M. Ikeda-Saito and S. Shaik, *J. Am. Chem. Soc.*, 2008, **130**, 14778–14790.
- 15 C. Lu, X. Zhao, Y. Lu, D. L. Rousseau and S.-R. Yeh, *J Am Chem Soc*, 2010, **132**, 1598–1605.
- 16 M. L. Quillin, T. Li, J. S. Olson, G. N. Phillips, Y. Dou, M. Ikeda-Saito, R. Regan, M. Carlson, Q. H. Gibson and H. Li, *J Mol Biol*, 1995, **245**, 416–436.

[View Online](#)